

Citrus Greening Disease (Huanglongbing)

Candidatus Liberibacter asiaticus
Ca. L. africanus
Ca. L. americanus

Primary hosts

Citrus sp. and some related plants, box thorn or Chinese box orange (*Severinia buxifolia*), wood apple (*Limonia acidissima*), white ironwood (*Vepris lanceolata*), and mock orange or orange jasmine (*Murraya paniculata*).

Symptoms

Symptoms are many and variable: yellow shoots, twig dieback, leaf drop, leaves with blotchy yellow/green coloration similar to the symptoms of zinc nutritional deficiency, enlarged veins that appear corky, excessive fruit drop, small and misshapen fruit, fruit that remains green at one end (the styler end) after maturity, fruit with mottled yellow/green coloration, small dark aborted seed inside fruit, discolored vascular bundles in the pithy center of the fruit, bitter tasting fruit, and silver spots left on fruits that are firmly pressed. (see <http://www.apsnet.org/online/feature/huanglongbing/> for symptom photographs).

The time from infection to the appearance of symptoms is variable, depending on the time of year, environmental conditions, tree age, host species/cultivar and horticultural health ranging from less than one year to several years.

The three disease agents (*Candidatus Liberibacter* spp.) are not distinguishable from each other based on symptoms produced.

Life cycle

*Candidatus Liberibacter*s are gram-negative bacteria with a double-membrane cell envelope. *Ca. L. asiaticus*, *africanus* and *americanus* are found in plants only in the phloem cells. The bacteria are transmitted by psyllids, a type of insect, as they feed. *Candidatus L. asiaticus* and *Candidatus L. americanus* are transmitted by the adults of the citrus psyllid *Diaphorina citri* Kuwayana. *Candidatus L. africanus* is transmitted by the adult psyllid *Trioza erytreae* Del Guercio. The bacteria can be acquired by the insects in the nymphal stages and the bacteria may be transmitted throughout the lifespan of the psyllid.

Eggs are laid on newly emerging leaves and hatch in 2-4 days. Five nymphal instars complete development in 11-15 days. The entire life cycle takes 15-47 days, depending upon temperature, and adults may live several months with females laying up to 800 eggs in a lifetime.

In an orchard, diseased trees are clustered together, with secondary infections produced 25 – 50m away. *Ca. L. africanus* is found at elevations greater than 700m and is less heat tolerant than *Ca. L. asiaticus*. *Ca. L. americanus* resembles *Ca. L. africanus*

in being less heat tolerant. Infections of *Ca. L. asiaticus* and *Ca. L. americanus* are more severe than *Ca. L. africanus* and can lead to tree death.

Current Geographic Distribution

Ca. L. africanus is found in eastern, central and southern Africa. *Ca. L. americanus* is found in Sao Paulo State, Brazil. *Ca. L. asiaticus* is found in Asia from Japan to S. China, SE Asia and the Indian subcontinent to Pakistan, the Arabian peninsula (not including Iran), Brazil, Cuba (2009), Dominican Republic (2009), Mexico (2009) and Florida (2005), and Louisiana (June 2008) in the U.S. The vector *D. citri*, is more widely spread in south and central America, including Mexico (at least since 2004), and in the U.S. in Texas (2001), Louisiana (May 2008), Alabama (August 2008), Georgia (August 2008), Mississippi (August 2008), South Carolina (August 2008), and California (September 2008), posing a threat to the citrus industry in these areas.

Impact in Oregon

Negligible.

References

Aubert, B. (1987) *Trioza erythrae* del Guercio and *Diaphorina citri* Kuwayama (Homoptera; Psylloidea), the two vectors of citrus greening disease: biological aspects and possible control strategies. *Fruits* 42:149-162

Chavan, V.M. and A.S. Summanwar (1993) Population dynamics and aspects of the biology of citrus psylla, *Diaphorina citri* Kuw. In Maharashtra. pp. 286-290 In P. Moreno, J.V. da Graca and L.W. Timmer [eds] Proc 12th Conference of the International organization of Citrus Virologists. University of California, Riverside.

Citrus Research Board (2008). The insect: Asian Citrus Psyllid.

<http://www.californiacitrustrustthreat.com/asian-citrus-psyllid.html>

Gottwald, T.R. (2007) Citrus Huanglongbing: the pathogen and its impact. APSnet feature, September 2007 <http://www.apsnet.org/online/feature/huanglongbing/>

Halbert, S.E. and K.L. Manjunath (2004) Asian citrus psyllids (Sternorrhyncha: Psyllidae) and greening disease of citrus: a literature review and assessment of risk in Florida. *Florida Entomologist* 87:330-353

Manjunath, K.L, S.E. Halbert, C. Ramadugu, S. Webb and R.F. Lee (2008) Detection of '*Candidatus Liberibacter asiaticus*' in *Diaphorina citri* and its importance in the management of citrus huanglongbing in Florida. *Phytopathology* 98:387-396

Martínez, Y., Llauger, R., Batista, L., Luis, M., Iglesia, A., Collazo, C., Peña, I., Casín, J. C., Cueto, J., Tablada, L. M. (2009) First report of '*Candidatus Liberibacter asiaticus*' associated with Huanglongbing in Cuba. *Plant Pathology* 58:389.

Matos, L., Hilf, M. E., Camejo, J. (2009) First report of '*Candidatus Liberibacter asiaticus*' associated with citrus Huanglongbing in the Dominican Republic. *Plant Disease* 93: 668 .

Mead, F.W. (1977) The Asiatic citrus psyllid, *Diaphorina citri* Kuwayama (Homoptera: Psyllidae) Entomology Circular 180. Florida Department of Agriculture and Consumer Services, Division of Plant Industry. 4pp.

NAPPO (2008) Phytosanitary alert system. Official pest reports. Confirmation of Asian citrus psyllid in San Diego county, California, United States.

<http://www.pestalert.org/oprDetail.cfm?oprID=343&keyword=citrus%20greening>

NAPPO (2008) Phytosanitary alert system. Official pest reports. Confirmation of Asian citrus psyllid, *Diaphorina citri* in Alabama, Georgia, Mississippi and South Carolina.

<http://www.pestalert.org/oprDetail.cfm?oprID=340&keyword=citrus%20greening>

NAPPO (2008) Phytosanitary alert system. Official pest reports. Confirmation of huanglongbing or citrus greening ('*Candidatus Liberibacter asiaticus*') in Louisiana, United States.

<http://www.pestalert.org/oprDetail.cfm?oprID=321&keyword=citrus%20greening>

NAPPO (2008) Phytosanitary alert system. Official pest reports. Detection of Asian citrus psyllid, *Diaphorina citri* Kuwayama, in Louisiana.

<http://www.pestalert.org/oprDetail.cfm?oprID=319&keyword=citrus%20greening>

NAPPO (2009) Phytosanitary alert system. Official pest reports. Detection of Huanglongbing ('*Candidatus Liberibacter asiaticus*') in the municipality of Tizimin, Yucatan, Mexico.

<http://www.pestalert.org/oprDetail.cfm?oprID=384>

Texeira, D.C., C. Saillard., S. Eveillard, J.L. Danet, P.I. da Costa, A.J. Ayres and J. Bove (2005) *Candidatus Liberibacter americanus*, associated with citrus huanglongbing (greening disease) in Sao Paulo State, Brazil. *Int. J. Syst. Evol. Microbiol* 55:1857-1862